

Virginia Save Our Streams <u>Monitor's Checklist for the Modified</u> Method



- 1) Choose a site (riffle) that is accessible (public property or with landowner permission) and that has the stream water bubbling over cobblestone sized rocks (3"-10" at the widest part of the particle). We strongly encourage monitors to avoid DEQ monitoring sites and the mixing zone of permitted wastewater discharges.
- 2) Use a Va. SOS seine net. This mesh is important for quality assurance purposes.
- 3) Approach the riffle from downstream (so as not to disturb potential collection areas) and position the net just below a spot with maximum bubbling action and a predominant number of cobbles. (approx. 45 degree angle) The net should be spread as widely as possible and set to allow a direct flow of water into the center of the net.
- 4) The sampling area is 1 square foot in front of the net. The net is 3 square feet, so you can use the size of the net to approximate your 1 square foot area. The monitor that will do the rubbing should take some cobbles from OUTSIDE the area to be sampled and rub them underwater (and outside of the "net zone") before gently laying them on the bottom of the net to anchor the net to the stream bottom.
- 5) The person holding the net will then time the other monitor to allow the rubbing of rocks for twenty seconds immediately upstream of the net. The final five seconds will be announced and for that time the "rubber" will scratch the stream bottom with their fingers or a garden cultivator type tool to collect any organism that live in the substrate. [20 seconds is ideal for the first time monitoring a new station, as you may end up with a lot of bugs to sort through. If you know the site and know that it does not yield many bugs in 20 seconds, you can increase the collection time up to 90 seconds, with 3/4 spent rubbing rocks and 1/4 spent disturbing the bottom].
- 6) Rub the "anchor" stones to remove any critters that may have attached themselves and with a forward and scooping motion remove the net from the stream. Examine the net for any organisms that are not macroinvertabrates (minnows or salamanders) and return them to the stream.
- 7) Take the net to the streamside and place it on a sheet that will allow for identification of any organisms that may pass through the mesh. Use ice cube trays and dishes to pick ALL organisms. Examine both sides of the net and the sheet beneath to obtain a rigorous count of all aquatic macroinvertabrates that were caught.
- 8) Repeat this procedure until a composite of all nets yields a total of organisms in excess of 200. Remember to thoroughly pick each net and add the total to the previous total. The time devoted to rubbing can be modified according to the judgment of the monitors but can not exceed 90 seconds per "dip". Also, no more than 4 "dips" can be made in pursuit of exceeding 200 organisms. If the monitors fail to find 200 organisms in 4 "dips" the calculation shall be made with the total that is obtained. Special note of this fact should be made in reporting the data.
- 9) With the individual counts of the organisms according to the categories as listed on the Va. SOS identification sheet and the total of all categories, calculate the six percentages (metrics) and combine them into one index value using the Va. SOS field calculation sheets. Be sure to report your results to Va. SOS ASAP.

Do this four times a year (every 3 months). If that is not possible, please focus on a spring and a fall collection at a minimum. Thank you for being a Virginia Save Our Streams monitor!

SAFETY

Things to remember when monitoring your stream...

- 1. Always remember to wash your hands after getting into any stream. The VA SOS method can not detect bacteriological pollution.
- 2. Glass may be hidden in the bottom of the stream watch out for it!
- 3. If you do get a cut or scrape while in the stream, use peroxide to clean the wound. Again, bacteriological pollution...
- 4. Always sample in pairs!
- 5. Do not sample a stream if water is above your knees or if water is flowing extremely fast. This can be dangerous.
- 6. Always obtain landowner permission before entering the stream.

POLLUTION

Sources of Pollution

When people talk about water, they talk about *point source pollution* and *nonpoint source pollution*

- 1. Point source pollution comes from a specific source: a pipe, a ditch, a container. It has a beginning point and an end point. Here's an easy way to remember, you can point to the pipe that's causing the problem.
- 2. Nonpoint source pollution comes from many scattered sources. It occurs when water (runoff) moves across and under the ground (think rain storm). The runoff picks up natural and man-made pollutants as its moves across the land. Then the runoff deposits the pollutants at the bottom of the watershed, into streams, rivers, lakes, estuaries, and even underground aquifers. Can you point to the problem? You might be able to point to different sources but you can't tell if, when, or how the source is getting into the waterbody.

Types of Pollution

- 1. Toxic pollution, like DDT or other chemicals that cause organisms to die and can threaten human health. Toxic pollution can come from pipes or barrels (point source), but it can also come from runoff (nonpoint source).
- 2. Sediment pollution can clog our waterways, ruin habitat and clog the gills of organisms in the stream. Lack of vegetative cover and impervious surfaces both have an impact on sedimentation.
- 3. Nutrient pollution can cause plant life in a stream to overgrow; depleting oxygen and sometimes causing the temperature of the stream to get too high. Nutrients can come from fertilizers used in lawns and gardens and animal waste or human waste (nonpoint source or point source).
- 4. Bacteria pollution can cause human health problems usually gastrointestinal. Bacteria pollution comes from animal and human waste (nonpoint source or point source).

Virginia Save Our Streams Stream Quality Survey

For Office Use Only
Name of Reviewer
Date Reviewed
Data sent to
VA SOS Data Entry Date

The purpose of this form is to aid you in gathering and recording important data about the health of your stream. By keeping accurate and consistent records of your observations and data from your macroinvertebrate count, you can document changes in ecological condition. Please note, this method was designed and tested for conditions in the state of Virginia and may not be appropriate in other areas.

Date			
Stream	Sta	tion # of	participants
Group or individual			
Name of <u>certified</u> * monitor			
County	Latitude	Longitude	
Location (please be specific)			
Average stream width	ft Avero	age stream depth	in
Flow rate: High Normal	Low	Negligible	-
Weather last 72 hours			
Water Temperature	°F (Please specify	vif reporting tempero	ature in Celsius)
Collection Time:		Other comments	s:
Net 1:sec			
Net 2:sec			
Net 3:sec			
Net 4:sec			_

Please send data sheets to your regional coordinator or to <u>VA SOS, Izaak Watlon League of America</u>, 707 Conservation Lane, Gaithersburg, MD 20878. If you have any questions about the modified method or this particular collection, please call Leah Miller at 301-548-0150 x. 219 or e-mail vasos@iwla.org.

* Your data is most useful when you pass your certification. Please contact VA SOS to schedule your certification!

Flat Worms Leeches Crayfish Sowbugs Scuds Mayflies Dragonflies and Damselflies Hellgrammites, Fishflies, and Alderflies	Macroinvertebrate	Tally	Count
Leeches Crayfish Sowbugs Scuds Stoneflies Mayflies Hellgrammites, Fishflies, and Alderflies	Worms		
Leeches Crayfish Sowbugs Scuds Stoneflies Mayflies Hellgrammites, Fishflies, and Alderflies			
Leeches Crayfish Sowbugs Scuds Stoneflies Mayflies Hellgrammites, Fishflies, and Alderflies			
Leeches Crayfish Sowbugs Scuds Stoneflies Mayflies Hellgrammites, Fishflies, and Alderflies	Flat Worms		
Sowbugs Scuds Stoneflies Dragonflies and Damselflies Hellgrammites, Fishflies, and Alderflies			
Sowbugs Scuds Stoneflies Dragonflies and Damselflies Hellgrammites, Fishflies, and Alderflies			
Scuds Stoneflies Mayflies Dragonflies and Damselflies Hellgrammites, Fishflies, and Alderflies	Leeches		
Scuds Stoneflies Mayflies Dragonflies and Damselflies Hellgrammites, Fishflies, and Alderflies			
Scuds Stoneflies Mayflies Dragonflies and Damselflies Hellgrammites, Fishflies, and Alderflies			
Scuds Stoneflies Mayflies Dragonflies and Damselflies Hellgrammites, Fishflies, and Alderflies	Crayfish		
Scuds Stoneflies Mayflies Dragonflies and Damselflies Hellgrammites, Fishflies, and Alderflies	· (Ca)		
Scuds Stoneflies Mayflies Dragonflies and Damselflies Hellgrammites, Fishflies, and Alderflies			
Scuds Stoneflies Mayflies Dragonflies and Damselflies Hellgrammites, Fishflies, and Alderflies	(lie)		
Stoneflies Mayflies Dragonflies and Damselflies Hellgrammites, Fishflies, and Alderflies	Sowbugs		
Stoneflies Mayflies Dragonflies and Damselflies Hellgrammites, Fishflies, and Alderflies			
Stoneflies Mayflies Dragonflies and Damselflies Hellgrammites, Fishflies, and Alderflies	-7/11/1		
Mayflies Dragonflies and Damselflies Hellgrammites, Fishflies, and Alderflies	Scuds		
Mayflies Dragonflies and Damselflies Hellgrammites, Fishflies, and Alderflies	ATTRACTOR OF THE PARTY OF THE P		
Mayflies Dragonflies and Damselflies Hellgrammites, Fishflies, and Alderflies	- Mary 1		
Dragonflies and Damselflies Hellgrammites, Fishflies, and Alderflies	Stoneflies		
Dragonflies and Damselflies Hellgrammites, Fishflies, and Alderflies	* *		
Dragonflies and Damselflies Hellgrammites, Fishflies, and Alderflies			
Dragonflies and Damselflies Hellgrammites, Fishflies, and Alderflies	/ \ / \		
Hellgrammites, Fishflies, and Alderflies	Mayflies		
Hellgrammites, Fishflies, and Alderflies			
Hellgrammites, Fishflies, and Alderflies	X X		
Hellgrammites, Fishflies, and Alderflies	//\ //\ 🛝		
Alderflies	Dragonflies and Damselflies		
Alderflies			
Alderflies	1		
Alderflies	Hollgrammitos Eighflios and		
	Alderflies		
	2111116		
	minne		
"""" } } -			
* "			

Macroinvertebrate	Tally	Count
Common Netspinning Caddisfly		
Most Caddisfly		
Beetles		
Midges		
Blackflies		
DIAGNICS		
Most True Fly		
Gilled Snail		
Lunged Snail		
Clams		
Other Subsurface Invertebrates		
TOTAL NUMBER OF ORGANISMS	I IN SAMPLE	

Individual Metrics

Metric Number	Metric Organism Group	Number of metric organism		Total number of organisms in the sample		Percent (This is your value for this metric)
1	Mayflies + Stoneflies + Most Caddisflies		÷		Multiply by 100	%
2	Common Netspinners		÷		Multiply by 100	%
3	Lunged Snails		÷		Multiply by 100	%
4	Beetles		÷		Multiply by 100	%

Metric 5 - % Tolerant

Taxon	Number
Worms	
Flatworms	
Leeches	
Sowbugs	
Scuds	
Dragonflies and Damselflies	
Midges	
Black Flies	
Lunged Snails	
Clams	
Total Tolerant	
Total number of organisms in sample	
Total Tolerant divided by the total	
number of organisms in the sample	
Multiply by 100	
This is your Value for Metric 5	

Metric 6 - % Non-Insects

Taxon	Number
Worms	
Flatworms	
Leeches	
Crayfish	
Sowbugs	
Scuds	
Gilled Snails	
Lunged Snails	
Clams	
Total Non-Insects	
Total number of organisms in sample	
Total Non-Insects divided by the total	
number of organisms in the sample	
Multiply by 100	
This is your Value for this Metric 6	

EXAMPLE

Metric	Metric Organism Group	Number of metric		Total number		Percent
1	Mayflies + Stoneflies +	80	÷	204	X 100	39.2%
2	Common Netspinners	40	÷	204	X 100	19.6%
3	Lunged Snails	0	÷	204	X 100	0%
4	Beetles	9	÷	204	X 100	4.4%

METRIC 5 - % Tolerant

Taxon	Number
Worms	10
Flatworms	0
Leeches	0
Sowbugs	5
Scuds	0
Dragonflies and Damselflies	5
Midges	20
Black Flies	10
Lunged Snails	0
Clams	10
Total Tolerant	60
Total Tolerant divided by the total	204
number of organisms in the sample	
Multiply by 100 - This is your Value	29.4

Metric 6 - % Non-Insects

77011 10 0 70 1 1011 E1150015				
Taxon	Number			
Worms	10			
Flatworms	0			
Leeches	0			
Crayfish	5			
Sowbugs	5			
Scuds	0			
Gilled Snails	10			
Lunged Snails	0			
Clams	10			
Total Non-Insects	40			
Total Non-Insects divided by the total	204			
number of organisms in the sample				
Multiply by 100 - This is your Value	19.6			

Metric Number	Metric Organism	Your Metric Value	2	1	0	
1	% Mayflies + Stoneflies + Most	39.2	Greater than 32.2 X	16.1 - 32.2	Less than 16.1	
2	% Common Netspinners	19.6	Less than 19.7 X	19.7 - 34.5	Greater than 34.5	
3	% Lunged Snails	0	Less than 0.3 X	0.3 - 1.5	Greater than 1.5	
4	% Beetles	4.4	Greater than 6.4	3.2 - 6.4 X	Less than 3.2	
5	% Tolerant	29.4	Less than 46.7 X	46.7 - 61.5	Greater than 61.5	
6	% Non-Insects	19.6	Less than 5.4	5.4 - 20.8 X	Greater than 20.8	
	Sı	ubtotals:	Total # of 2s: 4	Total # of 1s: 2	Total # of Os: O	
Multiply by 2: Multiply by 1: Multiply by 0: 8 2 0						
	the 3 subtotals to get	1				

Ecological conditions cannot be determined at this time (Gray Zone) (8)
_____Unacceptable ecological condition (0 to 7)

Virginia Save Our Streams Multimetric Index

Write your metric value from the previous page in the 2nd column (Your Metric Value). Determine whether each metric should get a score of 2,1, or 0 - depending upon the range of your metric value. Put a check in the appropriate box for your metric value under 2,1, or 0. Count the total number of 2's, 1's, and 0's. Follow the multiplication at the bottom of the chart to determine your Save Our Streams Multimetric Index score and determine whether the site has acceptable or unacceptable ecological condition. Please note: If after 4 nettings there are 0 macroinvertebrates, the multimetric score for the sample will be 0. If this occurs, please contact VA SOS or VA DEQ immediately.

Metric Number	Metric Organism	Your Metric Value	2	1	0
1	% Mayflies + Stoneflies + Most Caddisflies		Greater than 32.2	16.1 - 32.2	Less than 16.1
2	% Common Netspinners		Less than 19.7	19.7 - 34.5	Greater than 34.5
3	% Lunged Snails		Less than 0.3	0.3 - 1.5	Greater than 1.5
4	% Beetles		Greater than 6.4	3.2 - 6.4	Less than 3.2
5	% Tolerant		Less than 46.7	46.7 - 61.5	Greater than 61.5
6	% Non-Insects		Less than 5.4	5.4 - 20.8	Greater than 20.8
			Total # of 2s:	Total # of 1s:	Total # of Os:
		-	Multiply by 2:	Multiply by 1:	Multiply by 0:
		Subtotals:			
Now add	the 3 subtotals to ge	t the Virginia Sav	e Our Streams Multime	tric Index score:_	
	 _ Ecological con 	ditions canno	ecological condit ot be determined ble ecological con	at this time	(Gray Zone) (8)

Please send data sheets to your regional coordinator or VA SOS, 707 Conservation Lane, Gaithersburg, MD 20878. If you have any questions about the modified method or this particular collection, please call 301-548-0150 x 219 or e-mail vasos@iwla.org.

	1	
Fish water quality indicators	Barriers to fish movement	Surface water appearance
□ scattered individuals	□ beaver dams	□ clear □ milky
□ scattered schools	□ man-made dams	□ clear, tea colored □ black
□ trout (pollution sensitive)	□ waterfalls (>1 ft.)	□ colored sheen (oily)
□ bass (somewhat sensitive)	□ other	□ foamy □ other
□ catfish (pollution tolerant)	□ none	□ muddy
□ carp (pollution tolerant)		□ gray
Stream bed deposit (bottom)	Odor:	Stability of steam bed:
gray orange/red	none none	Bed sinks beneath your feet in:
□ yellow □ black	□ musky	□ no spots
□ brown □ silt	oil oil	□ a few spots
□ sand	□ sewage	□ many spots
□ other	other other	
Algae color:	Algae located:	Stream Channel Shade:
□ light green	□ everywhere	□ >75% full
□ dark green	□ in spots	□ 50%-74% high
□ brown coated	% bed covered	□ 25%-49% moderate
□ matted on stream bed	70 000 00 10100	□ 1%-24% slight
hairy		none
Stream bank composition		Riffle composition (=100%)
% trees	Stream bank erosion potential	% silt (mud)
% shrubs	□ >75% severe	% sand (1/64"-1/4" grains)
% grass	□ 50%-75% high	% gravel (1/4"-2" stones)
% bare soil	□ 25%-49% moderate	% cobbles (2"-10" stones)
% rocks	□ 1% - 24% slight	% boulders (>10" stones)
% other	□ none	/0 bounders (>10 stones)
	1 11 1 1 1 1 1 1	. 1 1
Land uses in the watershed : Record all land uses observed in the watershed area upstream and surrounding your sampling site. Indicate whether the following land uses have a high (H), moderate		
	_	- · · · · · · · · · · · · · · · · · · ·
	each the quality of your steam. (League to your watershad) Refer to	
procedures to determine how to a	esent in your watershed.) Refer to	the SOS standard operating
	1	
Oil & gas drilling	Sanitary landfill	Trash dump
Housing developments	Active construction	Fields
Forest	Mining (types)	Livestock pasture
Logging		Other
Urban uses (parking lots,	Cropland (types)	
highways, etc.		
Describe the amount of litter in and around the stream. Also describe the type of litter in and around the		
stream.		
Comments: Indicate what you think are the current and potential threats to your stream's health. Feel free to		
attach additional pages or photographs to better describe the condition of your		
stream.		